


Atomic structure

3.1.1.1 Fundamental particles

3.1.1.2 Mass number and isotopes

3.1.1.1 Fundamental particles

The atom is made up of a very dense, small and positively charged nucleus surrounded by negatively charged electrons. The electrons occur at specific energy levels, the principal energy levels are known as shells. The nucleus has neutrons and positively charged protons having a relative mass of 1 whereas the mass of an electron is so small about $1/1836$.



1. Electrons are released by the electron gun knock electrons off particles to produce + ions.
2. A negative plate attracts and accelerates the positive ions so they all have the same energy.
3. + ions pass through flight tube, as they all have the same energy lighter isotopes move faster.
4. At the detector, which is negatively charged, the ions are discharged producing a current.

3.1.1.2 Mass number and isotopes

Atomic mass (mass number) is number of protons + neutrons
Atomic number (proton number) is the number of protons in nucleus

Isotopes are atoms of the same element having the same atomic number but different mass numbers due to the difference in the number of neutrons. Isotopes have the same chemical properties but different physical properties.

3.1.1.2 Mass number and isotopes

Isotopes can be written down using the following notation:

A_ZX where A is the atomic mass and Z represents the atomic number. **Examples:** isotopes of hydrogen

| Name | Isotope | Number of neutrons |
|-----------|------------------|--------------------|
| Protium | ${}^1_1\text{H}$ | 0 |
| Deuterium | ${}^2_1\text{H}$ | 1 |
| Tritium | ${}^3_1\text{H}$ | 2 |

There are two methods of ionisation:

- **Electrospray ionisation** used for large molecules, less likely to fragment molecule. Sample is dissolved in water or alcohol, passed through hyperdermic with needle at high voltage so particle gains a proton: $X_{(g)} + \text{H}^+ \rightarrow \text{XH}^+_{(g)}$
- **Electron impact** is used for smaller molecules and will usually knock one electron per particle to produce positive ions: $X_{(g)} \rightarrow X^+_{(g)} + e^-$

Calculating relative atomic mass:

Abundance is the percentage of one isotope in a given sample of a mixture of isotopes.

$$\text{relative atomic mass } A_r = \frac{\text{combined mass of all isotopes}}{\text{combined abundance of all isotopes}}$$

Example: Calculate the RAM of silver which has the isotopes ${}^{107}\text{Ag}$ 51.88% and ${}^{109}\text{Ag}$ 48.12%.

$$A_r = \frac{107 \times 51.88 + 109 \times 48.12}{100} = 108$$

Exercise 1

1. Atoms of the same element having same number of protons but different number of neutrons.
2.
 - a. Both methods produce positive ions.
 - b. The needle in the electrospray method is at high voltage so causes particles to gain a proton. In the impact method electrons released by an electron gun will knock electrons from particles.

Exercise 2

1. When accelerated all isotopes are given the same energy so lighter isotopes will move faster than heavier isotopes.
2. Positive ions hit the detector, which is a negatively charged plate, the ions become discharged and produce a small current that is proportional to their abundance.
3.
 - a) $[{}^{35}\text{Cl} - {}^{37}\text{Cl}]^+$
 - b) $[{}^{37}\text{Cl} - {}^{37}\text{Cl}]^+$
4. relative atomic mass $A_r = \frac{\text{combined mass of all isotopes}}{\text{combined abundance of all isotopes}}$
5. $A_r = \frac{10.0 \times 19.8 + 11.0 \times 80.2}{100} = 10.8$
6. $A_r = \frac{28.0 \times 98.21 + 29.0 \times 4.70 + 30.0 \times 3.09}{100} = 29.8$